



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2009

MRI and clinical syndrome in dura materrelated Creutzfeldt-Jakob disease

Meissner, B ; Kallenberg, K ; Sanchez-Juan, P ; Ramljak, S ; Krasnianski, A ; Heinemann, U ;
Eigenbrod, S ; Gelpi, E ; Barsic, B ; Kretzschmar, H A ; Schulz-Schaeffer, W J ; Knauth, M ; Zerr, I

Abstract: **OBJECTIVE:** Iatrogenic Creutzfeldt-Jakob disease (iCJD) is mainly associated with dura mater (DM) grafts and administration of human growth hormones (hGH). Data on disease course in DM-CJD are limited. We describe the clinical and diagnostic findings in this patient group with special emphasis on MRI signal alterations. **METHODS:** Ten DM-CJD patients were studied for their clinical symptoms and diagnostic findings. The MRIs were evaluated for signal increase of the cortical and subcortical structures. **RESULTS:** DM-CJD patients had a median incubation time of 18 years and median disease duration of 7 months. The majority of patients were MM homozygous at codon 129 of the prion protein gene (PRNP) and presented with gait ataxia and psychiatric symptoms. No correlation between the graft site and the initial disease course was found. The MRI showed cortical and basal ganglia signal increase each in eight out of ten patients and thalamic hyperintensity in five out of ten cases. Of interest, patients with thalamic signal increase were homozygous for methionine. **CONCLUSION:** The MRI findings in DM-CJD largely resemble those seen in sporadic CJD, as the cortex and basal ganglia are mainly affected.

DOI: <https://doi.org/10.1007/s00415-009-0026-z>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-19661>

Journal Article

Published Version

Originally published at:

Meissner, B; Kallenberg, K; Sanchez-Juan, P; Ramljak, S; Krasnianski, A; Heinemann, U; Eigenbrod, S; Gelpi, E; Barsic, B; Kretzschmar, H A; Schulz-Schaeffer, W J; Knauth, M; Zerr, I (2009). MRI and clinical syndrome in dura materrelated Creutzfeldt-Jakob disease. *Journal of Neurology*, 256(3):355-363. DOI: <https://doi.org/10.1007/s00415-009-0026-z>

B. Meissner
K. Kallenberg
P. Sanchez-Juan
S. Ramljak
A. Krasnianski
U. Heinemann
S. Eigenbrod
E. Gelpi
B. Barsic
H. A. Kretzschmar
W. J. Schulz-Schaeffer
M. Knauth
I. Zerr

MRI and clinical syndrome in dura mater-related Creutzfeldt-Jakob disease

Received: 9 September 2007
Received in revised form: 29 April 2008
Accepted: 4 June 2008
Published online: 23 January 2009

B. Meissner, MD (✉) · A. Krasnianski, MD ·
U. Heinemann, MD · I. Zerr, MD
National TSE Reference Center
Dept. of Neurology
University of Göttingen
Robert-Koch-Str. 40
37075 Göttingen, Germany
Tel.: +49-551/39-6636
Fax: +49-551/39-7020
E-Mail: epicjd@med.uni-goettingen.de

present address:

B. Meissner, MD
Dept. of Gerontopsychiatry
University of Zürich
Minervastr. 145
8008 Zuerich, Switzerland

K. Kallenberg, MD · M. Knauth, MD
Dept. of Neuroradiology and MR research
in Neurology and Psychiatry
Göttingen, Germany

P. Sanchez-Juan, MD, PhD
Foundation Marqués de Valdecilla, IFIMAV
Santander, Spain

and

Centro de Investigación Biomédica en Red
sobre Enfermedades Neurodegenerativas
(CIBERNED)

S. Ramljak · W. J. Schulz-Schaeffer, MD
Dept. of Neuropathology
Georg-August University
Göttingen, Germany

S. Eigenbrod, MD · H. A. Kretzschmar, MD
Center for Neuropathology and Prion
Research
Ludwig-Maximilians-Universität
München, Germany

E. Gelpi, MD
Institute of Neurology
Medical University of Vienna
Vienna, Austria

B. Barsic, MD
University Hospital of Infectious Diseases
Zagreb, Croatia

■ **Abstract** *Objective* Iatrogenic Creutzfeldt-Jakob disease (iCJD) is mainly associated with dura mater (DM) grafts and administration of human growth hormones (hGH). Data on disease course in DM-CJD are limited. We describe the clinical and diagnostic findings in this patient group with special emphasis on MRI signal alterations. *Methods* Ten DM-CJD patients were studied for their clinical symptoms and

diagnostic findings. The MRIs were evaluated for signal increase of the cortical and subcortical structures. *Results* DM-CJD patients had a median incubation time of 18 years and median disease duration of 7 months. The majority of patients were MM homozygous at codon 129 of the prion protein gene (*PRNP*) and presented with gait ataxia and psychiatric symptoms. No correlation between the graft site and the initial disease course was found. The MRI showed cortical and basal ganglia signal increase each in eight out of ten patients and thalamic hyperintensity in five out of ten cases. Of interest, patients with thalamic signal increase were homozygous for methionine. *Conclusion* The MRI findings in DM-CJD largely resemble those seen in sporadic CJD, as the cortex and basal ganglia are mainly affected.

■ **Key words** CJD · MRI · dementia · cerebellar disorder · CNS infection

Introduction

Studies in iatrogenic CJD are extremely useful to understand some aspects of infections in humans, as the infec-

tious agent may reach the brain via different routes: direct inoculation (EEG depth electrodes), oral infection (growth hormone therapy, vCJD), peripheral infection (blood transfusion) and superficial infection (dura mater and corneal transplant).

The diagnosis of iCJD is based on the presence of typical symptoms (dementia, ataxia, extrapyramidal and pyramidal signs, myoclonus and akinetic mutism) and a history of iatrogenic exposure. The finding of 14-3-3 protein in the CSF may support the diagnosis (77 % sensitivity) [1, 2].

Dura mater grafts have been found as disease cause in 200 CJD patients worldwide [1]. Due to the long incubation periods ranging from 16 months to 31 years, DM-CJD cases are still detected nowadays [1, 3–9].

Increased rates of early ataxia have been reported in DM-CJD, hGH-CJD and vCJD [10–13]. In DM-CJD, a possible influence of the graft location on the clinical onset has been suggested [14, 15]. Only limited data were reported on paraclinical tests in iCJD, such as MRI. Since the MRI pattern of high signal abnormalities follows the lesion pattern of the brain as detected by pathology, the analysis of MRI in iCJD might be extremely useful.

The goals of our study were to describe MRI and clinical findings in a larger number of DM-CJD patients and to correlate them with the graft location and codon 129 genotype.

Methods

Patients with suspected CJD are reported to the CJD Surveillance Unit in Goettingen, (established in 1993) and examined by a study physician at the hospital reporting the case. A questionnaire about the patient's history (profession, habits, meat consumption) is filled out and copies are made of the medical charts (clinical findings, laboratory tests, EEG and MRI). The patients are classified as *possible*, *probable* or *no CJD* case according to established criteria [16]. One Croatian case (case 10) was included in this study because material was sent for genetic analysis and all clinical data were available.

Genetic analysis

Analysis of the prion protein gene (*PRNP*) was performed after isolation of genomic DNA from blood according to standard methods [17]. *PRNP* mutations were excluded in all but one patient by full *PRNP* sequencing (case 9).

Neuropathology

Histological examination was performed according to standard methods [18, 20].

Clinical examination

The patients were examined for the presence of typical CJD symptoms at onset (within first month) and during the disease course. Symptoms representing a residual state after surgery were not included in the analysis. Residual symptoms which progressed at the time of the disease onset were considered to be CJD signs and included in the study.

MRI

MRIs were available in nine out of ten patients. All MR examinations had been obtained from a 1.5 Tesla scanner (a total of 11 T2w, 9 FLAIR (fluid attenuated inversion recovery), 6 PDw (proton density weighted imaging) and 4 DWI (diffusion-weighted imaging), including serial MRIs in four patients) and were mainly available as hardcopy. Digital MRI data were available in four patients (cases 2, 3, 5 and 8). The MRI scans were reported independently and in consensus by two neuroradiologists (KK and MK) aware of the CJD diagnosis but not aware of the disease aetiology. The interobserver agreement was moderate for all the sequences (Concordance 80, and $\kappa = 0.52$) and high for FLAIR (Concordance 83.6 and $\kappa = 0.57$), PDw (Concordance 83.3 and $\kappa = 0.67$) and DWI (Concordance 83.9 and $\kappa = 0.65$). The results of the consensus review were used for the study.

Using a standardized protocol, *seven cortical areas* (the frontal, parietal, temporal, occipital and insula cortex, as well as the cingulate gyrus and hippocampus), the *basal ganglia* (caudate nucleus, putamen, globus pallidus), the *thalamus* and the *cerebellum* were assessed for signal increase in relation to suspected normointense tissue.

Results

Patients with dura mater grafts

In the years 1993 to 2006, ten DM-CJD cases (four *definite* and six *probable*) were identified (nine in Germany, one in Croatia) (Table 1). All surgeries were performed between 1980 and 1987 (Fig. 1). The median age at onset was 55 years (range 25–70); the median disease duration was 8 months (range 2–30). Incubation periods ranged from 9 to 23 years (median 18 years) (Fig. 1).

The codon 129 genotype was available in all patients (8 MM, 1 VV and 1 MV). Codon 219 was not examined in any case. The type of pathological prion protein (PrP^{Sc} 1 or 2) was analyzed in three out of four autopsied cases (only formalin fixed brain tissue was available in one patient): Three patients displayed PrP^{Sc} type 1. In one further case, the PrP^{Sc} type could not be clearly identified as type one or two (case 7) (as previously discussed) [21] (Table 1).

Clinical presentation

The main symptoms at onset (within the first month) and during the course of the disease are displayed in Table 2. Seven out of ten patients presented with gait ataxia (in three cases in isolation). Psychiatric abnormalities (depression and aggression) occurred in four out of ten patients (in one case as the only presenting symptom). Visual signs (double vision) were found in three patients, pyramidal signs (hemiparesis) in two and extrapyramidal signs (rigidity and choreatic movements) in one out of ten patients.


Only one patient displayed early signs of dementia. As unspecific clinical signs, vertigo (five patients) and headaches (five patients, data not displayed) were found most frequently. Further symptoms observed in the

Table 1 Clinical and diagnostic data in ten dura-related iatrogenic CJD cases

Case	Sex/age (years)	Codon 129 genotype (PrP ^{Sc} type)	Cause for surgery	CJD diagnosis (WHO criteria)	Incubation (years)	Disease duration (months)	EEG: PSWC	CSF: 14-3-3
1	m/38	MV	Brain trauma	Probable	10	4	+	+
2	m/44	MM (1)	Brain trauma	Definite	19	25	+	+
3	f/70	MM	Meningeoma	Probable	19	3	–	+
4	m/42	MM	AV-angioma	Probable	13	4	+	+
5	m/54	MM	Meningioma	Probable	18	11	–	+
6	f/56	MM (1)	Acousticus neurinoma	Definite	9	6	+	+
7	m/57	MM	Hemangioblastoma	Definite	20	10	–	+
8	f/65	MM	Trigeminal neuralgia	Probable	15	12	–	+
9	f/64	MM (1)	Pilocytic astrocytoma	Definite	23	2	+	+
10	f/25	VV	Pilocytic astrocytoma	Probable	17	30	(+)	(–)
Median 55 years					Median 18	8	5/9	9/9

M male; f female; PSWC periodic sharp wave complexes

(+), (–) = assessed as positive or negative at the hospital reporting the case (not included in the present analysis)

Fig. 1 Incubation periods (years) and disease duration (months) in patients with dura mater graft

early disease stage were dysaesthesias in three patients and loss of hearing, tinnitus, dysarthria, aphasia, abnormal sense of taste in one patient each.

Time course of symptoms

The median duration from the beginning of the disease to the onset of typical CJD symptoms is given in Fig. 2. Ataxia and psychiatric signs were among the earliest symptoms and followed by visual signs (median 1 month) and dementia (median 2 months). Myoclonus, extrapyramidal and pyramidal signs were typical signs of the advanced disease stage (median 3 to 4 months). Akinetic mutism was found in only two patients.

Dura graft location and initial clinical symptoms

Details on the graft placement and the clinical signs are shown in Table 2.

Early ataxia occurred most frequently in patients with suboccipital graft (four cases) but also in three patients with different graft location. Psychiatric signs were predominant in patients with frontal graft (three cases) but also found in one patient with suboccipital graft. Two patients with early pyramidal signs (hemiparesis) had a dura graft in the parietal area. Early rigidity was found in one patient with frontobasal graft. In four patients, the early disease course appeared to be monosymptomatic: three patients with suboccipital graft showed isolated ataxia for a time period of 4 weeks (case 9), 6 weeks (case 7) and 8 weeks (case 10). In one


Table 2 Dura graft location and clinical signs at onset and during course of the disease

Case	Dura graft location	Onset (1 st month of the disease)										Course**				
		Dementia	Ataxia	Visual	EPM	Pyramidal*	Other	Dementia	Ataxia	Visual	EPM	Pyramidal*	Myoclonus			
1	Fronto-basal	+	+	-	+	-	Psychiatric dysarthria	+	+	-	+	+	+			
2	Fronto-basal	-	-	-	-	-	Psychiatric	+	+	+	-	-	+			
3	Fronto-parietal	-	-	+	-	+	Psychiatric, aphasia	-	+	-	-	+	+			
4	Temporal	-	+	+	-	-	Dysaesthesias, vertigo	+	+	-	+	+	+			
5	Parieto-occipital	-	+	-	-	+	Dysaesthesias, vertigo	+	+	-	-	-	+			
6	Suboccipital	-	+	-	-	-	Psychiatric, dysaesthesias, loss of hearing and sense of taste	-	+	-	-	+	+			
7	Suboccipital	-	+	-	-	-	Vertigo	+	+	+	+	+	+			
8	Suboccipital	-	-	+	-	-	Vertigo, hyperhidrosis	+	+	+	+	+	+			
9	Suboccipital	-	+	-	-	-	-	+	+	+	+	+	+			
10	Suboccipital	-	+	-	-	-	Vertigo, tinnitus	+	+	-	+	-	+			
	-	1/10	7/10	3/10	1/10	2/10		8/10	10/10	4/10	6/10	7/10	10/10			

* corresponding to hemiparesis (two patients at onset) or spasticity (seven patients during the course)

** During the disease course all but one patient (case 5) developed psychiatric signs. Only two patients (cases 6 and 8) became akinetic and mute

EPM extrapyramidal signs

**Fig. 2** Time course of symptoms in patients with dura mater graft

further patient with frontobasal graft, only psychiatric signs were present for a time period of 6 months (case 2).

MRI

A synopsis of the MRI findings and applied sequences is given in Table 3. The cingulate gyrus was the most frequently affected cortex region (eight out of ten), followed by the frontal, temporal and parietal lobes (six out of ten, each). The occipital cortex was least affected (three out of ten). Basal ganglia hyperintensity was found in eight out of ten patients (in one case only during follow-up), and thalamic hyperintensity in five (in three cases only on PDw). In one patient (MV type), no signal abnormalities were found (Fig. 3). The current criteria for a pulvinar- or hockey stick sign (pulvinar or dorsomedial thalamus brighter than the other grey matter) were not met in any of the patients [22, 23].

Two patients showed pronounced signal increase in the brain hemisphere containing the graft: In one patient with right temporal graft, the signal increase was more pronounced in the right temporal and parietal lobes (Case 4). In one patient with left suboccipital graft, the left frontal cortex appeared to be more hyperintense than on the right side (Case 10). In two other patients, the findings were reversed (signal more pronounced in the opposite brain hemisphere) (Cases 6 and 9).

Discussion

In a recent study on DM-CJD cases, MRI sensitivity was reported to be high but no details were given on the involved brain areas [24]. According to a literature review, basal ganglia signal increase was found in eight out of 27 patients [6, 8, 9, 21, 25, 26], additional cortical signal increase in three [9, 25, 26] and thalamic hyperintensity in one [9]. In a large number of further cases, only atrophy was reported, which is most likely due to the less com-

Table 3 MRI signal increase in eight iatrogenic CJD cases caused by dura mater transplant

Case	Dura graft location	Genotype	Frontal	Temporal	Parietal	Occipital	Insula	Gc	Hc	Bg	Thalamus	Cerebellum	MRI 1 (time point)*	MRI 2 (time point)*
1	Fronto-basal	MV	-	-	-	-	-	-	-	-	-	-	T2, FLAIR (1/3)	-
2	Fronto-basal	MM	+	-/+	-/+	-	+	-	-	-/+	-	-	FLAIR, DWI (2/3)	T2, FLAIR (2/3)
3	Fronto-parietal	MM	-	+	-	+	+	+	-	+	+	-	T2, FLAIR (1/3)	-
4	Temporal	MM	+	+	+	+/-	-/+	+	+	+	-	-	T2, FLAIR (1/3)	T2, FLAIR, DWI (3/3)
5	Parieto-occipital	MM	+	+	+	+	+	+	+	+	+	-/+	T2, PD, DWI (2/3)	T2, PD, FLAIR, DWI (2/3)
6	Suboccipital	MM	+	-/+	+	+	+	+	-/+	+	-/+	-	T2, PD (2/3)	T2, PD, FLAIR (3/3)
7	Suboccipital	MM	-	-	-	-	-	-	-	+	+	+	PD (2/3)	-
8	Suboccipital	MM	+	+	+	-	+	+	-	+	+	-	T2, PD, DWI (1/3)	-
9	Suboccipital	MM	-	-	-	-	+	+	-	+	-	-	T2, FLAIR (2/3)	-
10	Suboccipital	VV	+	-	+	-	+	+	-	-	-	-	T2, FLAIR (1/3)	-
		-	6/10	6/10	6/10	3/10	5/10	8/10	3/10	8/10	5/10	2/10		

-/+ = signal increase during follow-up; +/- = signal increase not visible on follow-up MRI; Bg Basal ganglia; Gc Cingulate gyrus; Hc Hippocampus; * Time point given in thirds of the disease course




Fig. 3 Axial FLAIR (a) and diffusion-weighted (b) MRI of a 55-year old male patient (Case 5) six months after the disease onset showing signal increase in the basal ganglia, thalamus and cortex

mon use of FLAIR and DW images in the past (Table 4) [3, 5–9, 21, 25–41].

We studied the MRI findings in ten iatrogenic CJD patients after dura mater transplant by highly sensitive sequences. Cortex involvement was found in all but two patients. The basal ganglia were affected in eight, the thalamus in five and the cerebellum in two cases. The majority of our patients was MM homozygous at codon 129 of the *PRNP* and presented with gait ataxia.

At first sight, the MRI findings in our patients seem to largely resemble those seen in sporadic CJD. In sCJD, six molecular types (codon 129 genotype + PrP^{Sc} type 1 or 2; MM1, MM2, MV1, MV2, VV1, VV2) have been described with various clinical and diagnostic findings [42, 43]. Basal ganglia and cortical signal increase are most frequently found on the MRI [44–48]. Thalamic signal increase represents a comparatively rare finding, which seems to largely depend on the application of DW images [45–48].

Although the majority of our patients were not distinguishable from sporadic cases by means of MRI, the finding of thalamic hyperintensity in five out of eight MM homozygous patients is notable. According to previous literature reports, thalamic hyperintensity appears to be rather typical of MV2 type sporadic CJD and atypical of MM homozygous sCJD patients [49–51]. However, as only small patient groups have been studied so far with limited numbers of DW images, it remains to be confirmed that thalamic hyperintensity is rather not typical of MM homozygous sporadic CJD.

From the clinical point of view, sporadic CJD cases with MM homozygosity may present with rapidly progressive dementia (MM1 individuals), rather slowly developing cognitive signs (MM2-cortical individuals) or early ataxia (MM2-thalamic individuals with variable disease onset). The latter patients, who *clinically* most resemble our iatrogenic patients, are however well dis-

Table 4 Review of the literature: MRI findings and codon 129 genotype in dura-related iatrogenic CJD

Author	Gender/Age at onset (years)	Codon 129 genotype (PrP ^{Sc} type)	Dura mater introduction	Incubation (years)	Disease Duration (months)	MRI signal increase	Atrophy (CT/MRI)	MRI sequence	MRI performed (m.a.o.)
Kopp et al. 1996 [31]	f/52	MM (1)	Left sylvian artery aneurism 1984	11	5	–	–	n.m.	n.m.
Hannah et al. 2001 [30]	f/39	MM (1)	Arnold Chiari I malformation 1992	6	4	–	–	n.m.	n.m.
Nishida et al. 2002 [34]	f/69	MM (1)	Hemifacial spasm 1984 (Janetta's operation)	14	6	–	–	n.m.	n.m.
Mochizuki et al. 2003 [7]	f/79	MM (1)	Parasagittal meningioma 1982	15	18	–	+	T1, T2	n.m.
Wakisaka et al. 2006 [9]	f/19	MM (1)	Brain trauma 1985	17	10	Basal ganglia, pulvinar, left fronto-temporal cortex	–	T2, PD, DWI	1 week prior to death
Preusser et al. 2006 [8]	m/28	MM (2)	Brain trauma 1982	23	6	Basal ganglia	–	T2	n.m.
Delisle et al. 1993 [29]	m/18	MM	Craniopharyngeoma 1981, hGH 1982	8	12	–	+	n.m.	n.m.
Yamada et al. 1994 [41]	f/31	MM	Pituitary adenoma 1985	5	27	–	+	T1	12
Martínez-Lage et al. 1994 [33]	m/17	MM	Cerebellar astrocytoma 1983	7	3	–	+	T2	2nd week after admission or later
Martínez-Lage et al. 1994 [33]	m/25	MM	Arnold Chiari malformation 1983	9	15	–	+	T2	2nd week after admission or later
Yamada, et al. 1997 [40]	m/52	MM	Right frontal convexity meningioma 1984	8	10	White matter	+	T2	9
Takahima et al. 1997 [37]	f/38	MM	Aneurysm of right middle cerebral artery 1985	9	17	–	–	n.d.	–
Shimizu et al. 1999 [34]	m/68	MM	Hemifacial spasm 1985	11	9	–	+	n.m.	1
Shimizu et al. 1999 [36]	f/68	MM	Right parasagittal meningioma 1986	10	12	–	+	n.m.	n.m.
Kimura et al. 2001 [5]	m/42	MM	Pituitary adenoma 1984	14	13	–	–	DWI and other	n.m.
Groes et al. 2002 [28]	m/51	MM	Arteriovenous malformation 1983	14	6	–	–	n.m.	n.m.
Groes et al. 2002 [28]	m/44	MM	Hemangioblastoma 1988	10	4	–	–	n.m.	n.m.
Kobayashi et al. 2003 [25]	f/37	MM	Cerebellar astrocytoma (at the age of 14)	23	n.m.	Basal ganglia, Cortex	+	DWI	n.m.
Kretschmar et al. 2003 [21]*	m/58	MM	Angioblastoma 1980	19	18	Basal ganglia	–	n.m.	n.m.
Pocchiari et al. 1992 [35]	f/32	VV	Ethmoidal meningocoele 1981	10	8	–	+	n.m.	n.m.
Liscic et al. 1999 [32]*	m/24	VV	Brain trauma 1986	12	10	–	+	T2	8
Boutoleau et al. 2003 [3]	m/35	VV	Brain trauma 1991	10	2	–	–	n.m.	n.m.
Antoine et al. 1997 [27]	m/25	MV	Dura mater embolization in the external carotid artery for a nasopharyngeal angiofibroma	8	8	–	–	n.m.	n.m.
Thadani et al. 1988 [39]	f/28	n.m.	Cholesteatoma 1985	1.6	3	–	–	n.m.	n.m.
Takayama et al. 1993 [38]	m/38	n.m.	Cranial nerve decompression for hemifacial spasm 1985	5	n.m.	–	+	n.m.	n.m.
Lang et al. 2001 [6]	m/57	n.m.	Cerebellar angioblastoma 1980	19	19	Basal ganglia	–	DWI	n.m.
Ukisu et al. 2005 [26]	f/57	n.m.	n.m.	n.m.	n.m.	Basal ganglia, Cortex	–	T2, DWI	2 and 3 months

– = no abnormalities reported; n.m. not mentioned; m.a.o. months after onset; n.d. not applicable; * patients included in our study

Table 5 Disease phenotypes in sporadic and iatrogenic CJD with MM genotype

	Sporadic CJD			Iatrogenic CJD	
	Classic variant	Atypical variants with MM genotype		Dura mater-associated	vCJD
Genotype + PrP ^{Sc} type 1 or 2	MM1 ^a	MM2 – cortical ^b	MM2 – thalamic ^c	MM1 ^d	MM1 ^e
Infectious mode	–	–	–	Superficial	Oral
Age (median)	68 years (31–89)	67 years (60–82)	61 years (30–71)	56 years (19–79)	30 years (mean) (15–52)
Clinical onset	Dementia	Dementia	Variable	Ataxia	Ataxia
MRI signal increase	Cortex and basal ganglia	Cortex	None	Cortex, basal ganglia, thalamus	Cortex, basal ganglia, thalamus

^a [52] Collins et al. 2006; ^b [50] Krasnianski et al. 2006, [51] Hamaguchi et al. 2005; ^c 7 cases reported [51] Hamaguchi et al. 2005, [44] Shiga et al. 2004, [53] Yamashita et al. 2001; ^d 5 literature cases ([31] Kopp et al. 1996, [30] Hannah et al. 2001, [34] Nishida et al. 2002, [33] Mochizuki et al. 2003, [9] Wakisaka et al. 2006) and 3 own cases; ^e [54] Zeidler et al. 2000, [22] Collie et al. 2003

tinguished by MRI, as signal abnormalities are usually not observed in this particularly rare disease type. In MM2-cortical individuals, signal increase was mainly found in the cortex (Table 5) [7, 9, 22, 30, 31, 34, 44, 50–54].

Variant CJD patients, who may also present with early psychiatric signs and gait ataxia, may be differentiated from our cases by the presence of the *pulvinar sign* on the MRI. In the early disease stages, this finding may however be missing and follow-up exams should thus be performed [22]. Finally, a typical history of surgery may shift the diagnosis to iatrogenic CJD rather than sporadic or variant (Table 5) [7, 9, 22, 30, 31, 34, 44, 50–54].

Three of our MM homozygous patients did not show thalamic signal increase on the MRI. This might be explained by the MR examination time point and applied MR sequences, as thalamic signal changes were mainly seen in the later disease stages and on PDw images. Although the PDw sensitivity for thalamic hyperintensity has been reported to be high, it should be considered that information on the specificity of this sequence is very limited [22, 47].

Thalamic hyperintensity has also been reported in a Japanese dura recipient (MM type), four growth hormone recipients (two MV types, one VV type and one with unknown genotype) and one case of vCJD after blood transfusion (MM type) [9, 55, 56]. In conclusion, we have to consider the possibility that thalamic hyperintensity might be a peculiarity of acquired CJD forms.

In line with other studies, our patients were mainly MM homozygous at codon 129 of the *PRNP* [1, 11, 12, 26]. The prevalence of MM or VV homozygosity in iCJD has been discussed as due to increased susceptibility.

Methionine and valine homozygosity are associated with shorter incubation periods in iCJD [1, 57–59]. However, in DM-CJD cases, no such influence was found [60]. Compared to other reports, the incubation periods observed in our patients were significantly prolonged

(mean 18 years vs. 8 years), which may be best explained by the fact that German CJD surveillance was only first established in 1993 [11]. Iatrogenic CJD cases with shorter incubation periods were thus probably missed.

No clear correlation between the original graft location, the clinical syndrome and the MRI signal alterations was found, which may be due to varying examination time points and techniques or may indicate the independence of MRI signal alterations in iatrogenic CJD. Another possibility to discuss is that the clinical syndrome and MRI changes might be functions of the infectious agent's characteristics rather than the site of inoculation.

DM-CJD cases have recently been divided into a *plaque type* and *non-plaque type* on the basis of clinico-pathological findings. The plaque type usually presents with ataxia, whereas classic CJD features are found in the non-plaque type [24]. MRI hyperintensities were frequent in both types (not described in detail). In our study group, neuropathological data were limited and only one patient with florid plaques was found. This case presented with isolated ataxia, and the MRI showed hyperintensities of the cortex, basal ganglia and thalamus.

Summarizing, the MRI findings in DM-CJD are similar to sporadic CJD, showing frequent cortex and basal ganglia involvement. Thalamic hyperintensities in MM homozygotes might however be suggestive of an iatrogenic CJD form, rather than sporadic.

■ **Conflict of interest** The authors declare no conflict of interest.

■ **Acknowledgements** The authors thank Ms Bodemer, Ms Ciesielczyk and Mr Guettlich for technical assistance, Mr Giess for material supply and Ms Ehrlich and Ms Crozier for the manuscript processing. This study was funded by the Robert Koch Institute through funds from the Federal Ministry of Health (grant no. 1369-341) and by the (BMBF 01GI0301 and KZ: 0312720).

References

- Brown P, Brandel JP, Preese M, Sato T (2006) Iatrogenic Creutzfeldt-Jakob disease: the waning of an era. *Neurology* 67:389–393
- Sanchez-Juan P, Green A, Ladogana A, Cuadrado-Corralles N, Sanchez-Valle R, Mitrova E, Stoeck K, Sklavadias T, Kulczycki J, Hess K, Bodemer M, Slivarichova D, Saiz A, Calero M, Ingrosso L, Knight R, Janssens C, Van Duijn C, Zerr I (2006) Cerebrospinal fluid tests in the differential diagnosis of CJD. *Neurology* 67:637–643
- Boutoleau C, Guillon B, Martinez F, Vercelletto M, Faure A, Feve JR (2003) Iatrogenic Creutzfeldt-Jakob disease subsequent to dural graft: persisting risk after 1987. *Eur J Neurol* 10: 521–523
- Croes EA, Jansen GH, Lemstra AW, Frijns CJ, van Gool WA, van Duijn CM (2001) The first two patients with dura mater associated Creutzfeldt-Jakob disease in the Netherlands. *J Neurol* 248:877–880
- Kimura K, Nonaka A, Tashiro H, Yaginuma M, Shimokawa R, Okeda R, Yamada M (2001) Atypical form of dural graft associated Creutzfeldt-Jakob disease: report of a postmortem case with review of the literature. *J Neurol Neurosurg Psychiatry* 70: 696–699
- Lang CJ, Heckmann JG, Querner V, Neundorfer B, Kornhuber J, Buchfelder M, Kretschmar H (2001) Disease latency in Creutzfeldt-Jakob disease via dural grafting: a case report. *Eur J Epidemiol* 17:1013–1014
- Mochizuki Y, Mizutani T, Tajiri N, Oinuma T, Nemoto N, Kakimi S, Kiamoto T (2003) Creutzfeldt-Jakob disease with florid plaques after cadaveric dura mater graft. *Neuropathology* 23:136–140
- Preusser M, Strobel T, Gelpi E, Eiler M, Broessner G, Schmutzhard E, Budka H (2006) Alzheimer-type neuropathology in a 28 year old patient with iatrogenic Creutzfeldt-Jakob disease after dural grafting. *J Neurol Neurosurg Psychiatry* 77:413–416
- Wakisaka Y, Santa N, Doh-ura K, Kitamoto T, Ibayashi S, Iida M, Iwaki T (2006) Increased asymmetric pulvinar magnetic resonance imaging signals in Creutzfeldt-Jakob disease with florid plaques following a cadaveric dura mater graft. *Neuropathology* 26:82–88
- Fradkin JE, Schonberger LB, Mills JL, Gunn WJ, Piper JM, Wysocki DK, Thomson R, Durako S, Brown P (1991) Creutzfeldt-Jakob disease in pituitary growth hormone recipients in the United States. *JAMA* 265:880–884
- Hoshi K, Yoshino H, Urata J, Nakamura Y, Yanagawa H, Sato T (2000) Creutzfeldt-Jakob disease associated with cadaveric dura mater grafts in Japan. *Neurology* 55:718–721
- Lang CJ, Heckmann JG, Neundorfer B (1998) Creutzfeldt-Jakob disease via dural and corneal transplants. *J Neurol Sci* 160:128–139
- Will RG, Zeidler M, Stewart GE, Macleod MA, Ironside JW, Cousens SN, Mackenzie J, Estibeiro K, Green AJ, Knight RS (2000) Diagnosis of new variant Creutzfeldt-Jakob disease. *Ann Neurol* 47:575–582
- Iwasaki Y, Mimuro M, Yoshida M, Hashizume Y, Kitamoto T, Sobue G (2008) Clinicopathologic characteristics of five autopsied cases of dura mater-associated Creutzfeldt-Jakob disease. *Neuropathology* 28:51–61
- Heath CA, Barker RA, Esmonde TFG, Harvery P, Roberts R, Trend P, Head MW, Smith C, Bell JE, Ironside JW, Will RG, Knight RSG (2006) Dura mater-associated Creutzfeldt-Jakob disease: experience from surveillance in the UK. *J Neurol Neurosurg Psychiatry* 77:880–882
- WHO (1998) Human transmissible spongiform encephalopathies. *Wkly Epidemiol Rec* 47:361–365
- Windl O, Giese A, Schulz-Schaeffer W, Zerr I, Skworc K, Arendt S, Oberdieck C, Bodemer M, Poser S, Kretschmar HA (1999) Molecular genetics of human prion diseases in Germany. *Hum Genet* 105:244–252
- Kretschmar HA, Ironside JW, DeArmond SJ, Tateishi J (1996) Diagnostic criteria for sporadic Creutzfeldt-Jakob disease. *Arch Neurol* 53:913–920
- Parchi P, Castellani R, Capellari S, Ghetti B, Young K, Chen SG, Farlow M, Dickson DW, Sima AAF, Trojanowski JQ, Petersen RB, Gambetti P (1996) Molecular basis of phenotypic variability in sporadic Creutzfeldt-Jakob disease. *Ann Neurol* 39:767–778
- Schulz-Schaeffer WJ, Tschoke S, Kraneffuss N, Drose W, Hause-Reitner D, Giese A, Groschup MH, Kretschmar HA (2000) The paraffin-embedded tissue blot detects PrP(Sc) early in the incubation time in prion diseases. *Am J Pathol* 156:51–56
- Kretschmar H, Sethi S, Földvári Z, Windl O, Querner V, Zerr I, Poser S (2003) Iatrogenic Creutzfeldt-Jakob Disease with florid plaques. *Brain Pathol* 13:245–249
- Collie DA, Summers DM, Sellar RJ, Ironside JW, Cooper S, Zeidler M, Knight R, Will R (2003) Diagnosing variant Creutzfeldt-Jakob disease with the pulvinar sign: MR imaging findings in 86 neuropathologically confirmed cases. *Am J Neuroradiol* 24: 1560–1569
- WHO (2001) The Revision of the Surveillance Case Definition for Variant Creutzfeldt-Jakob Disease (vCJD): Report of a WHO Consultation Edinburgh, United Kingdom, 17 May 2001. Available at: <http://www.who.int/csr/resources/publications/bse/whocdscsreph20015.pdf>
- Noguchi-Shinohara M, Hamaguchi T, Kitamoto T, Sato T, Nakamura Y, Mizusawa H, Yamada M (2007) Clinical features and diagnosis of dura mater graft associated Creutzfeldt-Jakob disease. *Neurology* 69:360–367
- Kobayashi Y, Hirata K, Tanaka H, Yamada T (2003) Quinacrine administration to a patient with Creutzfeldt-Jakob disease who received a cadaveric dura mater graft – an EEG evaluation. *Rinsho Shinkeigaku* 43:403–408
- Ukisu R, Kushihashi T, Kitanosono T, Fujisawa H, Takenaka H, Ohgiya Y, Gokan T, Munechika H (2005) Serial diffusion-weighted MRI of Creutzfeldt-Jakob disease. *Am J Roentgenol* 184:560–566
- Antoine JC, Michel D, Bertholon P, Mosnier JF, Laplanche JL, Beaudry P, Hauw JJ, Veyret C (1997) Creutzfeldt-Jakob disease after extracranial dura mater embolization for a nasopharyngeal angiofibroma. *Neurology* 48: 1451–1453
- Croes EA, Roks G, Jansen GH, Nijssen PC, van Duijn CM (2002) Creutzfeldt-Jakob disease 38 years after diagnostic use of human growth hormone. *J Neurol Neurosurg Psychiatry* 72:792–793
- Delisle MB, Fabre N, Rochiccioli P, Doerr-Schott J, Rumeau JL, Bes A (1993) Creutzfeldt-Jakob disease after treatment with human extracted growth hormone. A clinicopathological study. *Rev Neurol Paris* 149: 524–527
- Hannah EL, Belay ED, Gambetti P, Krause G, Parchi P, Capellari S, Hoffman RE, Schonberger LB (2001) Creutzfeldt-Jakob disease after receipt of a previously unimplicated brand of dura mater graft. *Neurology* 56: 1080–1083
- Kopp N, Streichenberger N, Deslys JP, Laplanche JL, Chazot G (1996) Creutzfeldt-Jakob disease in a 52-year-old woman with florid plaques (letter). *Lancet* 348:1239–1240

32. Liscic RM, Brinar V, Miklic P, Barsic B, Himbele J (1999) Creutzfeldt-Jakob disease in a patient with a lyophilized dura mater graft. *Acta Med Croatica* 53:93–96
33. Martinez-Lage JF, Poza M, Sola J, Tortosa JG, Brown P, Cervenakova L, Esteban JA, Mendoza A (1994) Accidental transmission of Creutzfeldt-Jakob disease by dural cadaveric grafts. *J Neurol Neurosurg Psychiatry* 57: 1091–1094
34. Nishida Y, Yamada M, Hara K, Tsunemi T, Yamawaki M, Shimokawa R, Okeda R, Tsutsumi T, Mizusawa H (2002) Creutzfeldt-Jakob disease after Jannetta's operation with cadaveric dura mater graft: initial manifestations related to the grafted site. *J Neurol* 249:480–483
35. Pocchiari M, Masullo C, Salvatore M, Genuardi M, Galgani S (1992) Creutzfeldt-Jakob disease after non-commercial dura mater graft (letter). *Lancet* 340:614–615
36. Shimizu S, Hoshi K, Muramoto T, Homma M, Ironside JW, Kuzuhara S, Sato T, Yamamoto T, Kitamoto T (1999) Creutzfeldt-Jakob disease with florid-type plaques after cadaveric dura mater grafting. *Arch Neurol* 56: 357–362
37. Takashima S, Tateishi J, Taguchi Y, Inoue H (1997) Creutzfeldt-Jakob disease with florid plaques after cadaveric dural graft in a Japanese woman. *Lancet* 350:865–866
38. Takayama S, Hatsuda N, Matsumura K, Nakasu S, Handa J (1993) Creutzfeldt-Jakob disease transmitted by cadaveric dural graft: a case report. *No Shinkei Geka* 21:167–170
39. Thadani V, Penar PL, Partington J, Kalb R, Janssen R, Schonberger LB, Rabkin CS, Prichard JW (1988) Creutzfeldt-Jakob disease probably acquired from a cadaveric dura mater graft. Case report. *J Neurosurg* 69:766–769
40. Yamada M, Itoh Y, Suematsu N, Matsushita M, Otomo E (1997) Panencephalopathic type of Creutzfeldt-Jakob disease associated with cadaveric dura mater graft. *J Neurol Neurosurg Psychiatry* 63:524–527
41. Yamada S, Aiba T, Endo Y, Hara M, Kitamoto T, Tateishi J (1994) Creutzfeldt-Jakob disease transmitted by a cadaveric dura mater graft. *Neurosurgery* 34:740–743
42. Parchi P, Giese A, Capellari S, Brown P, Schulz-Schaeffer W, Windl O, Zerr I, Budka H, Kopp N, Piccardo P, Poser S, Rojiani A, Streichemberger N, Julien J, Vital C, Ghetti B, Gambetti P, Kretzschmar HA (1999) Classification of sporadic Creutzfeldt-Jakob disease based on molecular and phenotypic analysis of 300 subjects. *Ann Neurol* 46:224–233
43. Zerr I, Schulz-Schaeffer WJ, Giese A, Bodemer M, Schröter A, Henkel K, Tschampa HJ, Windl O, Pfahlberg A, Steinhoff BJ, Gefeller O, A. KH, Poser S (2000) Current clinical diagnosis in CJD: identification of uncommon variants. *Ann Neurol* 48:323–329
44. Shiga Y, Miyazawa K, Sato S, Fukushima R, Shibuya S, Sato Y, Konno H, Dohura K, Mugikura S, Tamura H, Higano S, Takahashi S, Itoyama Y (2004) Diffusion-weighted MRI abnormalities as an early diagnostic marker for Creutzfeldt-Jakob disease. *Neurology* 63:443–449
45. Young GS, Geschwind MD, Fischbein NJ, Martindale JL, Henry RG, Liu S, Lu Y, Wong S, Liu H, Miller BL, Dillon WP (2005) Diffusion-weighted and fluid-attenuated inversion recovery imaging in Creutzfeldt-Jakob disease: high sensitivity and specificity for diagnosis. *Am J Neuroradiol* 26:1551–1562
46. Tschampa HJ, Kallenberg K, Urbach H, Meissner B, Nicolay C, Kretzschmar HA, Knauth M, Zerr I (2005) MRI in the diagnosis of sporadic Creutzfeldt-Jakob disease: a study on inter-observer agreement. *Brain* 128:2026–2033
47. Kallenberg K, Schulz-Schaeffer WJ, Jastrow U, Poser S, Meissner B, Tschampa HJ, Zerr I, Knauth M (2006) Creutzfeldt-Jakob disease: Comparative Analysis of MR Imaging Sequences. *Am J Neuroradiol* 27: 1459–1462
48. Schroeter A, Zerr I, Henkel K, Tschampa HJ, Finkenstaedt M, Poser S (2000) Magnetic Resonance Imaging in the Clinical Diagnosis of Creutzfeldt-Jakob Disease. *Arch Neurol* 57:1751–1757
49. Krasnianski A, Schulz-Schaeffer WJ, Kallenberg K, Meissner B, Collie DA, Roeber S, Bartl M, Heinemann U, Vargas D, Kretzschmar HA, Zerr I (2006) Clinical findings and diagnostic tests in the MV-2 subtype of sporadic CJD. *Brain* 129:2288–2296
50. Krasnianski A, Meissner B, Schulz-Schaeffer W, Kallenberg K, Bartl M, Heinemann U, Vargas D, Kretzschmar H, Zerr I (2006) Clinical features and diagnosis of the MM2 cortical subtype of sporadic Creutzfeldt-Jakob disease. *Arch Neurol* 63:876–880
51. Hamaguchi T, Kitamoto T, Sato T, Mizusawa H, Nakamura Y, Noguchi M, Furukawa Y, Ishida C, Kuji I, Mitani K, Murayama S, Kohriyama T, Katayama S, Yamashita M, Yamamoto T, Uda F, Kawakami A, Ihara Y, Nishinaka T, Kuroda S, Suzuki N, Shiga Y, Arai H, Maruyama M, Yamada M (2005) Clinical diagnosis of MM2-type sporadic Creutzfeldt-Jakob disease. *Neurology* 64:643–648
52. Collins SJ, Sanchez-Juan P, Masters CL, Klug GM, van Duijn C, Poleggi A, Almonti S, Cuadrado-Corrales N, de Pedro-Cuesta J, Budka H, Gelpi E, Glatzel M, Seeger HE, Zerr I, Kretzschmar H, Jansen GH, Olsen E, Mitrova E, Alperovitch A, Brandel JP, Murray K, Will RG (2006) Determinants of diagnostic investigation results across the clinical spectrum of sporadic CJD. *Brain* 129:2278–2287
53. Yamashita M, Yamamoto T, Nishinaka K, Uda F, Kameyama M, Kitamoto T (2001) Severe brain atrophy in a case of thalamic variant of sporadic CJD with plaque-like PrP deposition. *Neuropathology* 21:138–143
54. Zeidler M, Ironside JW (2000) The new variant of Creutzfeldt-Jakob disease. *Rev Sci Tech* 19:98–120
55. Guray C, Zeghoudi AC, Atlaoui-Rabia S, Charitanski D, Mallecourt J (2005) Cerebral magnetic resonance imaging in iatrogenic Creutzfeldt-Jakob disease. *Presse Med* 34:1080–1081
56. Wroe SJ, Pal S, Siddique D, Hyare H, Macfarlane R, Joiner S, Linehan JM, Brandner S, Wadsworth JD, Hewitt P, Collinge J (2006) Clinical presentation and pre-mortem diagnosis of variant Creutzfeldt-Jakob disease associated with blood transfusion: a case report. *Lancet* 368:2061–2067
57. Brandel JP, Preece M, Brown P, Croes E, Laplanche JL, Agid Y, Will RG, Alperovitch A (2003) Distribution of codon 129 genotype in human growth hormone-treated CJD patients in France and the UK. *Lancet* 362:128–130
58. Deslys JP, Jaegly A, d'Aignaux JH, Mouthon F, de Villemeur TB, Dormont D (1998) Genotype at codon 129 and susceptibility to Creutzfeldt-Jakob disease (letter). *Lancet* 351:1251
59. Huillard d'Aignaux J, Costagliola D, Maccario J, Billelte de Villemeur T, Brandel JP, Deslys JP, Hauw JJ, Chausain JL, Agid Y, Dormond D, Alperovitch A (1999) Incubation period of Creutzfeldt-Jakob disease in human growth hormone recipients in France. *Neurology* 53:1197–1201
60. Brown P, Preece M, Brandel J-P, Sato T, McShane L, Zerr I, Fletcher A, Will RG, Pocchiari M, Cashmann NR, d'Aignaux JH, Cervenakova L, Fradkin J, Schonberger LB, Collins SJ (2000) Iatrogenic Creutzfeldt-Jakob disease at the millennium. *Neurology* 55:1075–1081